



Genomic and ecological variation in comammox Nitrospira populations

Palomo, Alejandro; Smets, Barth F.

Publication date:
2018

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Palomo, A., & Smets, B. F. (2018). *Genomic and ecological variation in comammox Nitrospira populations*. Poster session presented at 17th International Symposium on Microbial Ecology, Leipzig, Germany.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Genomic and ecological variation in comammox *Nitrospira* populations

Alejandro Palomo* and Barth F. Smets

Department of Environmental Engineering, Technical University of Denmark, Kgs Lyngby, Denmark.





Introduction and Objectives

- ✓ Complete ammonia oxidizing (comammox) bacteria belong to *Nitrospira*, an extremely diverse genus of nitrite oxidizing bacteria.
- ✓ Described comammox *Nitrospira* genomes belong to *Nitrospira* lineage II, and comprise two clades (clade A and B) based on the phylogeny of their ammonia monooxygenases.
- ✓ *Nitrospira* spp. are widespread in both natural and engineered ecosystems where nitrogen cycling occurs; multiple species frequently coexist.
- ✓ However, the forces driving differentiation and diversification in *Nitrospira* populations remain poorly understood.
- ✓ The purpose of this study was to investigate the distribution of comammox and other *Nitrospira* spp. across different environments and geographical locations.

Approach


- ✓ 55 different metagenome-assembled genomes (MAG) of *Nitrospira* were retrieved from 13 Danish groundwater-fed rapid sand filters and multiple publicly available metagenomes by combining different binning tools (MetaBAT¹, MaxBin2² and CONCOCT³) followed by binning refinement (Binning refiner⁴, DAS Tool⁵ and metaWRAP⁶) and dereplication (dRep⁷).
- ✓ These *Nitrospira* MAGs together with 37 published *Nitrospira* genomes were clustered (genomes with ANI $\geq 95\%$ were classified as member of same species) to create a database of *Nitrospira* species.
- ✓ The abundance of each *Nitrospira* species across 1000 metagenomes was assessed using MIDAS⁸.
- ✓ Distribution patterns of *Nitrospira* genomes across the metagenomes was visualized through principal component analysis (PCA).

Global distribution of *Nitrospira* spp.




More *Nitrospira* genomes were detected in metagenomes from DWTP (drinking water treatment plant) and freshwater samples compared to soil and WWTP (wastewater treatment plant) samples. Only DWTP metagenomes contained multiple abundant *Nitrospira* genomes (coverage > 5).

Genomic and diversity patterns of *Nitrospira* spp.




Comammox *Nitrospira* Clade A genomes (purple and pink) form two different clusters: CladeA1 genomes (purple) co-occur with non-comammox *Nitrospira* lineage I genomes and are present in WWTP samples; CladeA2 genomes (pink) are present in DWTP, groundwater and freshwater samples.

AmoA-based phylogeny



HaoA-based phylogeny



AmoA and HaoA sequences phylogenies are not congruent. While AmoA phylogeny separates comammox *Nitrospira* into two clades (A and B), HaoA phylogeny suggests existence of a subdivision within clade A (CladeA1 and CladeA2)

Conclusions

- ✓ *Nitrospira* genomes were detected in metagenomes from very different environments and geographical locations; a higher abundance was observed in DWTP metagenomes.
- ✓ Environment more than geography separates different *Nitrospira* spp.: WWTP metagenomes separate strongly from DWTP metagenomes based on presence and type of *Nitrospira* genomes.
- ✓ *Nitrospira* spp. affiliated to the same lineage or clade inhabit similar environments.
- ✓ Environmental partitioning within the comammox Clade A was detected. A similar separation was observed based on the phylogeny of hydroxylamine oxidoreductase subunit A amino acid sequences.

